COLORADO ASTHMA CARE PLAN AND MEDICATION ORDER FOR SCHOOL AND CHILD CARE SETTINGS (2018-2019)			
PARENT/GUARDIAN COMPLETE AN		ND SIGN: School/grade:	
Chil		, 8	
Pare	ent/Guardian	Name: Phone:	
Healthcare Provider Name:		Phone:	
Triggers: Weather (cold air, wind)		☐ Illness ☐ Exercise ☐ Smoke ☐ Dust ☐ Pollen ☐	7 Other:
Triggers: ☐ Weather (cold air, wind) ☐ Illness ☐ Exercise ☐ Smoke ☐ Dust ☐ Pollen ☐ Other: ☐ Life threatening allergy, specify:			
I give permission for school personnel to share this information, follow this plan, administer medication and care for my child/			
youth, and if necessary, contact our healthcare provider. I assume full responsibility for providing the school/program			
prescribed medication and supplies, and to comply with board policies, if applicable. I am aware 911 may be called if a quick			
relief inhaler is not at school and my child/youth is experiencing symptoms. I approve this care plan for my child/youth.			
PARENT SIGNATURE DATE NURSE/CCHC SIGNATURE DATE			
HEALTHCARE PROVIDER		QUICK RELIEF (RESCUE) MEDICATION: Albuterol Other:	
COMPLETE ALL ITEMS,		Common side effects: $\uparrow \uparrow$ heart rate, tremor \Box Have child use spacer with inhaler.	
		Controller medication used at home:	
IF YOU SEE THIS:		DO THIS:	
ii s	No current symptoms		outing Ctudent/Desert
GREEN ZONE: No Symptoms		Pretreat strenuous activity: ☐ Not required ☐ Routine ☐ Student/Parent request Give QUICK RELIEF MED 1015 minutes before activity: ☐ 2 puffs ☐ 4 puffs	
NZ	• Doing usual activities	Repeat in 4 hours, if needed for additional physical activity.	
If child is cu		child is currently experiencing symptoms, follow YELLOW ZONE.	
5 Z			TELLOW ZONE.
	Trouble breathing	1. Stop physical activity.	
ns ns		2. Give QUICK RELIEF MED: ☐ 2 puffs ☐ 4 puffs	
O to	• Frequent cough	3. Stay with child/youth and maintain sitting position.	
2 7	Complains of tight chest	4. REPEAT QUICK RELIEF MED, if not improving in 15 minutes: ☐ 2 puffs ☐ 4 puffs	
YELLOW ZONE: Mild symptoms	 Not able to do activities, 	5. Child/youth may go back to normal activities, once symptoms are relieved.	
	but talking in complete	6. Notify parents/guardians and school nurse.	
i) symptoms do not improve or worsen, johow RED ZONE.		D ZONE.	
	• Peak flow:&		
₹	 Coughs constantly 	1. Give QUICK RELIEF MED: ☐ 2 puffs ☐ 4 puffs	
EN ns	Struggles to breathe	Refer to anaphylaxis plan, if child/youth has lifethreatening allergy.	
RGI	 Trouble talking (only speaks 	2. Call 911 and inform EMS the reason for the call.	
VE	35 words)	3. Stay with child/youth. Remain calm, encouraging slower, deeper breaths.	
RED ZONE: EMERGENCY Severe Symptoms	Skin of chest and/or neck	4. Notify parents/guardians and school nurse.	
NE ere	pull in with breathing	5. If symptoms do not improve, REPEAT QUICK RELIEF MED: 2 puffs 4 puffs	
20	• Lips/fingernails gray or blue	every 5 minutes until EMS arrives.	
ED S	• ↓↓ Level of consciousness • Peak flow <	School personnel should not drive student to hospit	tai.
~	- Feak How <		
PROVIDER INSTRUCTIONS FOR QUICK RELIEF INHALER USE: CHECK APPROPRIATE BOX(ES)			
Light Student needs supervision or assistance to use inhaler. Student will not selfcarry inhaler.			
Light Student understands proper use of asthma medications, and in my opinion, can carry and use his/her inhaler at school			
independently with approval from school nurse and completion of contract.			
☐ Student will notify school staff after using quick relief inhaler, if symptoms do not improve with use.			
HEALTH CARE PROVIDER SIGNATURE PRINT PROVIDER NAME DATE FAX PHONE			
Copies of plan provided to: ☐Teacher(s) ☐PhysEd/Coach ☐Principal ☐Main Office ☐Bus Driver Other			

